THE FIRST AND ONLY FDA-APPROVED MECHLORETHAMINE GEL

VALCHLOR®
(mechlorethamine)gel
0.016%

Your Guide to Common Questions About MF-CTCL and VALCHLOR

What is VALCHLOR? PAGE 4

How does VALCHLOR work? PAGE 4

What is VALCHLOR® used for? ↓



Actor portrayal.

Please see glossary on PAGE 12 for definition of terms.

VALCHLOR is a prescription medicine used on the skin (topical) to treat people with Stage IA and IB mycosis fungoides-type cutaneous T-cell lymphoma (MF-CTCL) who have received previous skin treatment. It is not known if VALCHLOR is safe and effective in children.

IMPORTANT SAFETY INFORMATION

What is the most important information I should know about VALCHLOR?


VALCHLOR is for use on skin only. Keep VALCHLOR away from your eyes, mouth, and nose.

Please see Important Safety Information throughout and click here for the full Prescribing Information and Medication Guide for VALCHLOR.


What is MF-CTCL?

MF-CTCL stands for **mycosis fungoides-type cutaneous T-cell lymphoma**. It's a type of cancer that starts in white blood cells called T-cells. Normally, T-cells help your body fight infections. In MF-CTCL, some T-cells stay active all the time and multiply, causing buildup on your skin. That's why you may notice patches or plaques on your skin.


What MF-CTCL may look like



Stage IA–Patch



Stage IB–Plaque



Stage IA/IB–Hypopigmentation

Actual patient examples. Individual patient presentations may vary. Images used with permission from VisualDx.

Please see glossary on page 12 for definition of terms.

IMPORTANT SAFETY INFORMATION (CONT'D)

What is the most important information I should know about VALCHLOR? (cont'd)

VALCHLOR is for use on skin only. Keep VALCHLOR away from your eyes, mouth, and nose.

- If VALCHLOR gets in your eyes it can cause eye pain, burning, swelling, redness, sensitivity to light, and blurred vision. It may also cause blindness and permanent injury to your eyes.
- If VALCHLOR gets in your mouth or nose it can cause pain, redness, and ulcers.

In early stages (IA and IB), MF-CTCL stays in the skin

VALCHLOR® (mechlorethamine)gel 0.016%



Stage IA
Less than 10% of your skin is affected by patches or plaques.



Stage IB
10% or more of your skin is affected, but the cancer has not spread beyond the skin.

● Patches and/or plaques of MF-CTCL

For illustrative purposes only. Percentages and appearance may vary depending on a person's body type and size.

Early-stage MF-CTCL is most often treated with medicines applied directly to the skin. Oral medications and infusions are not usually needed at this stage. **VALCHLOR is made specifically to treat MF-CTCL Stages IA/IB.**

Scan or click here to learn more about MF-CTCL from the Cutaneous Lymphoma Foundation (CLF)

IMPORTANT SAFETY INFORMATION (CONT'D)

What is the most important information I should know about VALCHLOR? (cont'd)

Get medical help right away if VALCHLOR gets in your eyes, mouth, or nose. Talk with your healthcare provider before using VALCHLOR about how to get medical help.

Please see Important Safety Information throughout and click here for the full Prescribing Information and Medication Guide for VALCHLOR.

VALCHLOR CAN HELP

What is VALCHLOR?

VALCHLOR is a prescription gel used on the skin to treat Stage IA/IB MF-CTCL in people who have already tried another topical therapy. It is the **first and only FDA-approved** gel form of mechlorethamine.

How does VALCHLOR work?

The active ingredient in VALCHLOR is **mechlorethamine**. It works by destroying rapidly growing T-cells in your skin. Think of it as going right to the source—the abnormal cells in your skin—and stopping them from spreading.

How long has VALCHLOR been used?

For over a decade, VALCHLOR has been prescribed to help treat Stage IA/IB MF-CTCL.

Please see glossary on page 12 for definition of terms.

IMPORTANT SAFETY INFORMATION (CONT'D)

When should I not use VALCHLOR?

Do not use VALCHLOR if you are severely allergic to mechlorethamine. Tell your healthcare provider if you have had an allergic reaction to mechlorethamine.

Please see Important Safety Information throughout and click here for the full Prescribing Information and Medication Guide for VALCHLOR.


VALCHLOR® (mechlorethamine)gel 0.016%

How is VALCHLOR different?

You may have used steroid creams or had light therapy (phototherapy) before. VALCHLOR works in a different way. It is topical chemotherapy that destroys the rapidly growing T-cells. It works directly on the skin and **does not get absorbed into your bloodstream**.

You can apply VALCHLOR in the comfort of your own home.

Actor portrayal.



WATCH & LEARN

Scan or click here to see how to apply VALCHLOR

IMPORTANT SAFETY INFORMATION (CONT'D)

What should I tell my healthcare provider?

Before using VALCHLOR, tell your healthcare provider about all your medical conditions, including if you:

- are pregnant or plan to become pregnant. VALCHLOR may harm your unborn baby. Avoid becoming pregnant during treatment with VALCHLOR. Tell your healthcare provider right away if you think you are or may become pregnant during treatment with VALCHLOR.

When will I see a difference?

Treatment takes time. In a **yearlong** clinical study, some people noticed a difference as early as one month, and most others saw results in 6-12 months.

Every person's treatment journey with MF-CTCL is unique. It's important to apply as instructed by your healthcare provider. Skipping doses may delay results.



Actor portrayals.

Please see glossary on page 12 for definition of terms.

IMPORTANT SAFETY INFORMATION (CONT'D)

What should I tell my healthcare provider? (cont'd)

- **Females** who are able to become pregnant should use a barrier method of birth control, such as a male condom or spermicide, during treatment with VALCHLOR.
- **Males** with female partners who are able to become pregnant should use a barrier method of birth control, such as a male condom or spermicide, during treatment with VALCHLOR.

What is important to keep in mind when using VALCHLOR?

Talk with your healthcare team and follow their guidance for the best results. You can track your own progress by taking photos of your skin and sharing them with your healthcare provider (HCP). **Don't stop treatment early** without talking to your HCP first. Staying on your treatment may help improve outcomes.

VISIT VALCHLOR.COM

Or scan to get tips to help you stay on track with VALCHLOR

VALCHLOR was studied in the largest controlled clinical study of people with early-stage (IA/IB) MF-CTCL.

IMPORTANT SAFETY INFORMATION (CONT'D)

What should I tell my healthcare provider? (cont'd)

- are breastfeeding or plan to breastfeed. It is not known if VALCHLOR passes into your breast milk. Do not breastfeed during treatment with VALCHLOR.

  Tell your healthcare provider about all the medicines you take including prescription and over-the-counter medicines, vitamins, and herbal supplements.

Please see Important Safety Information throughout and click here for the full Prescribing Information and Medication Guide for VALCHLOR.

What side effects do I need to know?

VALCHLOR can cause serious side effects including risk of secondary exposure to VALCHLOR, inflammation of your skin (dermatitis), and increased risk of certain types of skin cancers.

Inflammation of the skin (dermatitis) is the most common side effect and may include redness, swelling, or itching at the application site. These reactions are common and, in some cases, may be manageable without discontinuing treatment; however, you should follow your HCP's guidance. **Get medical help right away if VALCHLOR gets in your eyes, mouth, or nose.**



Actor portrayal.

Please see glossary on page 12 for definition of terms.

IMPORTANT SAFETY INFORMATION (CONT'D)

What should I avoid while using VALCHLOR?

VALCHLOR is flammable. Avoid fire, flame, and smoking until VALCHLOR has dried.

Please see Important Safety Information throughout and click here for the full Prescribing Information and Medication Guide for VALCHLOR.

VALCHLOR® (mechlorethamine)gel 0.016%

What you may notice and should watch for when using VALCHLOR

Some skin changes are expected and manageable. Others need your doctor's attention right away. Here's a simple guide:

✔

Manage most common skin reactions with your care team:

- Redness or pinkness at the application site
- Itching or dryness

!

Call your HCP right away if you notice:

- Redness or swelling
- Ulcers or open sores
- Signs of skin infection
- Itching or blisters
- Darkening of areas of your skin

IMPORTANT SAFETY INFORMATION (CONT'D)

What are the possible side effects of VALCHLOR? (cont'd)

VALCHLOR can cause serious side effects, including:

- **Risk of secondary exposure to VALCHLOR.** You may have a risk of inflammation of your skin (dermatitis), injury to your eyes, mouth, or nose, and certain types of cancers. Caregivers who accidentally come into contact with VALCHLOR must wash the affected area thoroughly with soap and water right away for at least 15 minutes and remove any contaminated clothing.

Get medical help right away if VALCHLOR gets in your eyes, mouth, or nose.

What is VALCHLOR Support®?

VALCHLOR Support® is a program designed to help you start and stay on treatment, and is delivered through Accredo Specialty Pharmacy. Accredo handles everything from processing your prescription to delivering it right to your door. VALCHLOR is not available at your local retail pharmacy. VALCHLOR is **only** available through Accredo.

VALCHLOR Copay Program: eligible patients may pay no more than $0 per prescription fill*

How can I get my prescription?

VALCHLOR is delivered **directly to your home** through Accredo Health Group, a dedicated specialty pharmacy.

VISIT VALCHLOR.COM

Scan this code or click here to visit the website for step-by-step instructions for Accredo

Please see glossary on page 12 for definition of terms.

*The VALCHLOR copay program is only available for US and Puerto Rico residents who are 18 or older and have commercial health insurance with copay/coinsurance exceeding $0 per prescription fill. Patients ineligible for the VALCHLOR copay program include those enrolled in Medicare, Medicaid, VA/DoD (TRICARE), the Indian Health Service, or any other federal- or state-funded healthcare program, or where prohibited by law. The VALCHLOR copay program is not prescription drug coverage or insurance. Helsinn Therapeutics (U.S.), Inc. reserves the right to terminate or modify this program at any time or without notice. Other terms and conditions apply.

IMPORTANT SAFETY INFORMATION (CONT'D)

What are the possible side effects of VALCHLOR? (cont'd)

Get medical help right away if VALCHLOR gets in your eyes, mouth, or nose.

- **Inflammation of your skin (dermatitis)** is common with VALCHLOR and may sometimes be severe. Your risk for dermatitis is increased if VALCHLOR is applied to your face, genital area, anus, or skin folds. Tell your healthcare provider if you develop skin reactions such as redness, swelling, itching, blisters, ulcers, and skin infections.
- **Increased risk of certain types of skin cancers.** Certain types of skin cancer can develop on areas of your skin that are treated with VALCHLOR and areas of your skin that are not treated with VALCHLOR. Your healthcare provider will check your skin for skin cancers during and after your treatment with VALCHLOR. Tell your healthcare provider if you get any new skin lesions.

The most common side effects of VALCHLOR include: redness, swelling, itching, skin ulcers or blisters, skin infection, and darkening of areas of your skin.

VALCHLOR may cause fertility problems in females and males. This could affect the ability to become pregnant. Talk to your healthcare provider if this is a concern for you.

These are not all of the possible side effects of VALCHLOR. Call your doctor for medical advice about side effects. You may report side effects to FDA at **1-800-FDA-1088.**

Keep VALCHLOR and all medicines out of the reach of children.

Please see the full Prescribing Information and Medication Guide for VALCHLOR.

Scan or click here to download the Accredo specialty pharmacy app to help manage your VALCHLOR prescription

Please see Important Safety Information throughout and click here for the full Prescribing Information and Medication Guide for VALCHLOR.

Glossary

Bloodstream: The blood that flows throughout your body. Reminder–VALCHLOR works on the skin and does not get absorbed into your bloodstream.

Cutaneous: Relating to or affecting the skin.

Lesions: Any area of skin affected by MF-CTCL, including patches and plaques.

MF-CTCL: Mycosis fungoides T-cell lymphoma is a type of cutaneous T-cell lymphoma (CTCL) that causes patches or plaques to build up on the skin. MF-CTCL may also be called CTCL, cutaneous lymphoma, skin lymphoma, or mycosis fungoides-type CTCL.

Patches: Flat skin lesions that may appear red, pink, darker, lighter, dry, or scaly. They are often one of the earliest signs of MF-CTCL.

Plaques: Raised, thickened areas of skin that may be scaly or itchy.

Stage IA/IB MF-CTCL: MF-CTCL is staged to describe how much of your skin is affected by patches or plaques and whether the disease has spread beyond the skin or not. Stage IA covers less than 10% of the skin, while Stage IB covers 10% or more.

T-cells: A type of white blood cell that normally help your body fight infections. In MF-CTCL, some T-cells stay active and rapidly multiply.

Topical: A medicine you apply on the skin, rather than take as a pill or receive as an infusion.

Track your progress

VALCHLOR®
(mechlorethamine)gel
0.016%

Consider tracking any changes you notice, such as a reduction in itchiness or redness, or changes in the size of the patch or plaque. Be sure to discuss any changes with your doctor at your next visit.

MONTH

1

2

3

4

5

6

7

8

9

10

11

12

Please see Important Safety Information throughout and click here for the full Prescribing Information and Medication Guide for VALCHLOR.

Notes and questions for my doctor

VALCHLOR®
(mechlorethamine)gel
0.016%



Please see Important Safety Information throughout and click here for the full Prescribing Information and Medication Guide for VALCHLOR.

VALCHLOR®
(mechlorethamine)gel
0.016%

VALCHLOR key facts

- VALCHLOR is the first and only FDA-approved mechlorethamine gel made specifically to treat Stage IA and IB MF-CTCL
- VALCHLOR works directly on the skin and does not get absorbed into your bloodstream
- Treatment may take time. It's important to talk to your healthcare team about the length of treatment time and managing side effects
- Do not stop treatment with VALCHLOR before speaking with your HCP

Scan to learn more about VALCHLOR or visit VALCHLOR.com

What is VALCHLOR® used for?

VALCHLOR is a prescription medicine used on the skin (topical) to treat people with Stage IA and IB mycosis fungoides-type cutaneous T-cell lymphoma who have received previous skin treatment. It is not known if VALCHLOR is safe and effective in children.

SELECT IMPORTANT SAFETY INFORMATION

Do not use VALCHLOR if you are severely allergic to mechlorethamine. Tell your healthcare provider if you have had an allergic reaction to mechlorethamine.

Please see Important Safety Information throughout and click here for the full Prescribing Information and Medication Guide for VALCHLOR.

HELSINN

VALCHLOR® and VALCHLOR Support® are trademarks of Helsinn Birex Pharmaceuticals Ltd., Ireland. VALCHLOR® is distributed and marketed by Helsinn Therapeutics (U.S.), Inc. under license.
© 2026 Helsinn Therapeutics (U.S.), Inc.
All rights reserved. V-VALC-US-0519 08/26